

Fifth Metatarsal Fracture Rehabilitation Exercises

As soon as you can tolerate pressure on the ball of your foot, begin stretching your foot using the towel stretch. When this stretch is too easy, try the standing calf stretch and soleus stretch.

- **Towel stretch:** Sit on a hard surface with one leg stretched out in front of you. Loop a towel around your toes and the ball of your foot and pull the towel toward your body keeping your knee straight. Hold this position for 15 to 30 seconds then relax. Repeat 3 times.
- **Standing calf stretch:** Facing a wall, put your hands against the wall at about eye level. Keep one leg back with the heel on the floor, and the other leg forward. Turn your back foot slightly inward (as if you were pigeon-toed) as you slowly lean into the wall until you feel a stretch in the back of your calf. Hold for 15 to 30 seconds. Repeat 3 times and then switch the position of your legs and repeat the exercise 3 times. Do this exercise several times each day.
- **Standing soleus stretch:** Stand facing a wall with your hands on a wall at about chest level. With both knees slightly bent and one foot back, gently lean into the wall until you feel a stretch in your lower calf. Angle the toes of your back foot slightly inward and keep your heel down on the floor. Hold this for 15 to 30 seconds. Return to the starting position. Repeat 3 times.

You can do the next 5 exercises when your foot swelling has stopped increasing.

- **Ankle range of motion:** Sitting or lying down with your legs straight and your knee toward the ceiling, move your ankle up and down by pointing your toes toward your nose, then away from your body; in toward your other foot and out away from your other foot; and in circles. Only move your foot and ankle. Don't move your leg. Repeat 10 times in each direction. Push hard in all directions.



- **Resisted ankle dorsiflexion:** Sit with one leg out straight and your foot facing a doorway. Tie a loop in one end of elastic tubing. Put your foot through the loop so that the tubing goes around the arch of your foot. Tie a knot in the other end of the tubing and shut the knot in the door. Move backward until there is tension in the tubing. Keeping your knee straight, pull your foot toward your body, stretching the tubing. Slowly return to the starting position. Do 3 sets of 10.
- **Resisted ankle eversion:** Sit with both legs stretched out in front of you, with your feet about a shoulder's width apart. Tie a loop in one end of elastic tubing. Put one foot through the loop so that the tubing goes around the arch of that foot and wraps around the outside of the other foot. Hold onto the other end of the tubing with your hand to provide tension. Turn the foot with the tubing up and out. Make sure you keep your other foot still so that it will allow the tubing to stretch as you move your foot with the tubing. Return to the starting position. Do 3 sets of 10.

You may do the rest of the exercises when you can stand on your injured ankle without pain.

- **Heel raise:** Balance yourself while standing behind a chair or counter. Using the chair to help you, raise your body up onto your toes and hold for 5 seconds. Then slowly lower yourself down without holding onto the chair. Hold onto the chair or counter if you need to. When this exercise becomes less painful, try lowering on one leg only. Repeat 10 times. Do 3 sets of 10.
- **Standing toe raise:** Stand with your feet flat on the floor, rock back onto your heels and lift your toes off the floor. Hold this for 5 seconds. Do 3 sets of 10.
- **Towel pickup:** With your heel on the ground, pick up a towel with your toes. Release. Repeat 10 to 20 times. When this gets easy, add more resistance by placing a book or small weight on the towel.

- **Balance and reach exercises**

Stand upright next to a chair with your injured leg farthest from the chair. This will provide you with support if you need it. Stand just on the foot of your injured leg. Try to raise the arch of this foot while keeping your toes on the floor.

1. Keep your foot in this position and reach forward in front of you with the hand farthest away from the chair, allowing your knee to bend. Repeat this 10 times while maintaining the arch height. This exercise can be made more difficult by reaching farther in front of you. Do 2 sets.
2. Stand in the same position as above. While maintaining your arch height, reach the hand farthest away from the chair across your body toward the chair. The farther you reach, the more challenging the exercise. Do 2 sets of 10.


- **Single leg balance:** Stand without any support and attempt to balance on one leg. Begin with your eyes open and then try to perform the exercise with your eyes closed. Hold the single-leg position for 30 seconds. Repeat 3 times. When you have mastered this, try doing this exercise standing on a pillow.

Written by Tammy White, MS, PT, and Phyllis Clapis, PT, DHSc, OCS, for RelayHealth.

Published by RelayHealth.

© 2009 RelayHealth and/or its affiliates. All Rights Reserved

Fifth Metatarsal Fracture Rehabilitation Exercises



Towel stretch




Standing calf stretch



Standing soleus stretch



Ankle active range of motion



Fifth Metatarsal Fracture Rehabilitation Exercises



Heel raise



Standing toe raise



Towel pickup



Balance and reach exercise A



Balance and reach exercise B



Single leg balance