

Meniscal (Cartilage) Tear Rehabilitation Exercises

You may do the first 5 exercises right away. You may do the rest of the exercises when the pain in your knee has decreased.

- Passive knee extension: Do this exercise if you are unable to fully extend your knee. While lying on your back, place a rolled-up towel underneath the heel of your injured leg so the heel is about 6 inches off the ground. Relax your leg muscles and let gravity slowly straighten your knee. You may feel some discomfort while doing this exercise. Try to hold this position for 2 minutes. Repeat 3 times. Do this exercise several times per day. This exercise can also be done while sitting in a chair with your heel on another chair or stool.
- **Heel slide:** Sit on a firm surface with your legs straight in front of you. Slowly slide the heel of your injured leg toward your buttock by pulling your knee to your chest as you slide. Return to the starting position. Do 3 sets of 10.
- Standing calf stretch: Facing a wall, put your hands against the wall at about eye level. Keep one leg back with the heel on the floor, and the other leg forward. Turn your back foot slightly inward (as if you were pigeon-toed) as you slowly lean into the wall until you feel a stretch in the back of your calf. Hold for 15 to 30 seconds. Repeat 3 times and then switch the position of your legs and repeat the exercise 3 times. Do this exercise several times each day.
- Hamstring stretch on wall: Lie on your back with your buttocks close to a doorway, and extend your legs straight out in front of you along the floor. Raise one leg and rest it against the wall next to the door frame. Your other leg should extend through the doorway. You should feel a stretch in the back of your thigh. Hold this position for 15 to 30 seconds. Repeat 3 times and then switch legs and do the exercise again.
- Straight leg raise: Lie on your back with your legs straight out in front of you. Bend the knee on your uninjured side and place the foot flat on the floor. Tighten the thigh muscle of the other leg and lift it about 8 inches off the floor, keeping the thigh muscle tight throughout. Slowly lower your leg back down to the floor. Do 3 sets of 10.
- 281-256-8685
- 281-256-8879
- info@HealthZoneMD.com
- 607 Park Grove Drive Suite A, Katy, TX 77450



- Wall squat with a ball: Stand with your back, shoulders, and head against a wall and look straight ahead. Keep your shoulders relaxed and your feet 2 feet away from the wall and a shoulder's width apart. Place a soccer or basketball-sized ball behind your back. Keeping your back upright, slowly squat down to a 45-degree angle. Your thighs will not yet be parallel to the floor. Hold this position for 10 seconds and then slowly slide back up the wall. Repeat 10 times. Build up to 3 sets of 10.
- **Step-up:** Stand with the foot of your injured leg on a support (like a small step or block of wood) 3 to 5 inches high. Keep your other foot flat on the floor. Shift your weight onto your injured leg on the support straighten your knee as the other leg comes off the floor. Lower your leg back to the floor slowly. Do 3 sets of 10.
- **Knee stabilization:** Wrap a piece of elastic tubing around the ankle of the uninjured leg. Tie a knot in the other end of the tubing and close it in a door.
 - 1. Stand facing the door on the leg without tubing and bend your knee slightly, keeping your thigh muscles tight. While maintaining this position, move the leg with the tubing straight back behind you. Do 3 sets of 10.
 - 2. Turn 90 degrees so the leg without tubing is closest to the door. Move the leg with tubing away from your body. Do 3 sets of 10.
 - 3. Turn 90 degrees again so your back is to the door. Move the leg with tubing straight out in front of you. Do 3 sets of 10.
 - 4. Turn your body 90 degrees again so the leg with tubing is closest to the door. Move the leg with tubing across your body. Do 3 sets of 10.

Hold onto a chair if you need help balancing. This exercise can be made even more challenging by standing on a pillow while you move the leg with tubing.

- Resisted terminal knee extension: Make a loop from a piece of elastic tubing by tying a knot in both ends. Close both knots in a door. Step into the loop so the tubing is around the back of your injured leg. Lift the other foot off the ground. Hold onto a chair for balance, if needed. Bend the knee on the leg with tubing
- 281-256-8685
- **281-256-8879**
- info@HealthZoneMD.com
- 607 Park Grove Drive Suite A, Katy, TX 77450



about 45 degrees. Slowly straighten your leg, keeping your thigh muscle tight as you do this. Do this 10 times. Do 3 sets. An easier way to do this is to stand on both legs for better support while you do the exercise.

Written by Tammy White, MS, PT, and Phyllis Clapis, PT, DHSc, OCS, for RelayHealth. Published by RelayHealth.

© 2009 RelayHealth and/or its affiliates. All Rights Reserved.




281-256-8879

info@HealthZoneMD.com

0 607 Park Grove Drive Suite A, Katy, TX 77450



Meniscal (Cartilage) Tear Rehabilitation Exercises



607 Park Grove Drive Suite A, Katy, TX 77450