

Patellar Tendonitis (Jumper's Knee) Rehabilitation Exercises

You can do the first 4 exercises right away. When you have less pain in your knee, you can do the remaining exercises.

- **Standing hamstring stretch:** Place the heel of your injured leg on a stool about 15 inches high. Keep your knee straight. Lean forward, bending at the hips until you feel a mild stretch in the back of your thigh. Make sure you do not roll your shoulders and bend at the waist when doing this or you will stretch your lower back instead of your leg. Hold the stretch for 15 to 30 seconds. Repeat 3 times.
- **Quadriceps stretch:** Stand an arm's length away from the wall with your injured leg farthest from the wall. Facing straight ahead, brace yourself by keeping one hand against the wall. With your other hand, grasp the ankle of your injured leg and pull your heel toward your buttocks. Don't arch or twist your back. Keep your knees together. Hold this stretch for 15 to 30 seconds.
- **Side-lying leg lift:** Lying on your uninjured side, tighten the front thigh muscles on your top leg and lift that leg 8 to 10 inches away from the other leg. Keep the leg straight and lower slowly. Do 3 sets of 10.
- **Straight leg raise:** Lie on your back with your legs straight out in front of you. Bend the knee on your uninjured side and place the foot flat on the floor. Tighten the thigh muscle of the other leg and lift it about 8 inches off the floor, keeping the thigh muscle tight throughout. Slowly lower your leg back down to the floor. Do 3 sets of 10.
- **Step-up:** Stand with the foot of your injured leg on a support (like a small step or block of wood) 3 to 5 inches high. Keep your other foot flat on the floor. Shift your weight onto your injured leg on the support straighten your knee as the other leg comes off the floor. Lower your leg back to the floor slowly. Do 3 sets of 10.
- **Wall squat with a ball:** Stand with your back, shoulders, and head against a wall and look straight ahead. Keep your shoulders relaxed and your feet 2 feet away from the wall and a shoulder's width apart. Place a soccer or basketball-sized ball behind your back. Keeping your back upright, slowly squat down to a 45-degree angle. Your thighs will not yet be parallel to the floor. Hold this position for 10

 281-256-8685

 281-256-8879

 info@HealthZoneMD.com

 607 Park Grove Drive Suite A, Katy, TX 77450

seconds and then slowly slide back up the wall. Repeat 10 times. Build up to 3 sets of 10.

- **Knee stabilization:** Wrap a piece of elastic tubing around the ankle of the uninjured leg. Tie a knot in the other end of the tubing and close it in a door.
 1. Stand facing the door on the leg without tubing and bend your knee slightly, keeping your thigh muscles tight. While maintaining this position, move the leg with the tubing straight back behind you. Do 3 sets of 10.
 2. Turn 90 degrees so the leg without tubing is closest to the door. Move the leg with tubing away from your body. Do 3 sets of 10.
 3. Turn 90 degrees again so your back is to the door. Move the leg with tubing straight out in front of you. Do 3 sets of 10.
 4. Turn your body 90 degrees again so the leg with tubing is closest to the door. Move the leg with tubing across your body. Do 3 sets of 10.

Hold onto a chair if you need help balancing. This exercise can be made even more challenging by standing on a pillow while you move the leg with tubing.

- **Resisted terminal knee extension:** Make a loop from a piece of elastic tubing by tying a knot in both ends. Close both knots in a door. Step into the loop so the tubing is around the back of your injured leg. Lift the other foot off the ground. Hold onto a chair for balance, if needed. Bend the knee on the leg with tubing about 45 degrees. Slowly straighten your leg, keeping your thigh muscle tight as you do this. Do this 10 times. Do 3 sets. An easier way to do this is to stand on both legs for better support while you do the exercise.
- **Decline eccentric squat:** Stand with both feet on an angled platform or with your heels on a 3-inch-high board. Put all your weight on your injured leg and squat down to a 45-degree angle. Use your other leg to help you return to a standing position from the squat. You should lower your body to a squat using only your injured leg but you can return to an upright position using both legs. When this exercise gets easy, hold weights in your hands to make the exercise more difficult. Do 3 sets of 10.

Written by Tammy White, MS, PT, and Phyllis Clapis, PT, DHSc, OCS, for RelayHealth.



Published by RelayHealth.

© 2009 RelayHealth and/or its affiliates. All Rights Reserved.

-  281-256-8685
-  281-256-8879
-  info@HealthZoneMD.com
-  607 Park Grove Drive Suite A, Katy, TX 77450

Patellar Tendonitis (Jumper's Knee) Rehabilitation Exercises



Standing hamstring stretch



Quadriceps stretch



Side-lying leg lift




Straight leg raise



Patellar Tendonitis (Jumper's Knee) Rehabilitation Exercises



Knee stabilization: A



Knee stabilization: B



Knee stabilization: C

