

## Snapping Hip Syndrome Rehabilitation Exercises

You may do all of these exercises right away.

- **Quadriceps stretch:** Stand an arm's length away from the wall with your injured leg farthest from the wall. Facing straight ahead, brace yourself by keeping one hand against the wall. With your other hand, grasp the ankle of your injured leg and pull your heel toward your buttocks. Don't arch or twist your back. Keep your knees together. Hold this stretch for 15 to 30 seconds.
- **Hamstring stretch on wall:** Lie on your back with your buttocks close to a doorway, and extend your legs straight out in front of you along the floor. Raise one leg and rest it against the wall next to the door frame. Your other leg should extend through the doorway. You should feel a stretch in the back of your thigh. Hold this position for 15 to 30 seconds. Repeat 3 times and then switch legs and do the exercise again.
- **Gluteal stretch:** Lying on your back with both knees bent, rest the ankle of one leg over the knee of your other leg. Grasp the thigh of the bottom leg and pull that knee toward your chest. You will feel a stretch along the buttocks and possibly along the outside of your hip on the top leg. Hold this for 15 to 30 seconds. Repeat 3 times.
- **Iliotibial band stretch: Standing:** Cross one leg in front of the other leg and bend down and touch your toes. You can move your hands across the floor toward the front leg and you will feel more stretch on the outside of your thigh on the other side. Hold this position for 15 to 30 seconds. Return to the starting position. Repeat 3 times. Reverse the positions of your legs and repeat.
- **Iliotibial band stretch: Side-leaning:** Stand sideways near a wall. Place one hand on the wall for support. Cross the leg farthest from the wall over the other leg, keeping the foot closest to the wall flat on the floor. Lean your hips into the wall. Hold the stretch for 15 seconds, repeat 3 times, and then switch legs and repeat the exercise another 3 times.
- **Prone hip extension:** Lie on your stomach with your legs straight out behind you. Tighten the buttocks and thigh muscles of your injured leg and lift it off the floor about 8 inches. Keep your knee straight. Hold for 5 seconds. Then lower your leg and relax. Do 3 sets of 10.



281-256-8685



281-256-8879



[info@HealthZoneMD.com](mailto:info@HealthZoneMD.com)



607 Park Grove Drive Suite A, Katy, TX 77450



- **Side-lying leg lift:** Lying on your uninjured side, tighten the front thigh muscles on your top leg and lift that leg 8 to 10 inches away from the other leg. Keep the leg straight and lower slowly. Do 3 sets of 10.

Written by Tammy White, MS, PT, and Phyllis Clapis, PT, DHSc, OCS, for RelayHealth.

Published by RelayHealth.

© 2009 RelayHealth and/or its affiliates. All Rights Reserved.



281-256-8685



281-256-8879



[info@HealthZoneMD.com](mailto:info@HealthZoneMD.com)



607 Park Grove Drive Suite A, Katy, TX 77450

## Snapping Hip Syndrome Rehabilitation Exercises



Quadriceps stretch




Hamstring stretch on wall



Gluteal stretch



Iliotibial band stretch (standing)



Iliotibial band stretch (side-leaning)



Prone hip extension



Side-lying leg lift